

Wilchem's Sentinel Range has been formulated to protect micronutrients in solution to allow for the correction of nutrient deficiencies.

Стор Туре	Application	Rate Kg/Ha	Timing	Water (L/Ha)
Cereals, Canola, Cotton, Maize, Summer Crops	Foliar	0.26 to 1	Apply from mid tillering to flag leaf emergence	76
	Soil	1 to 3		
Lucerne, Legumes, Pasture	Foliar	1 to 1.5	10 to 14 days prior to flowering	76

POISON - Keep out of reach of children.

FIRST AID - Avoid contact with eyes and skin. If poisoning occurs contact a doctor or poisons information centre on 131 126.

Any recommendations are a guide and must be adapted to sult local conditions. Adjust rates based on tissue tests, soil tests, local knowledge and the recommendation of your agronomist. Do not exceed a solution of 0.1% on fruit crops and 1% on broadscre crops **Compatibility:** It is compatible with a wide range of crop protection products. It is stable in alkaline soils and in the presence of soluble phosphates. A jar test should be performed and/or test application before preparing a full tank mix.

Directions for use: Do not apply during periods of crop stress, extreme weather, flowering or fruit colour change. Ensure all other products are mixed in the tank before adding. Do not apply as foliar to stone fruit.



Batch Number