

MetaBOOST

Bio-active metabolites of Trichoderma spp. for rapid induction of the plant immune and stress response pathways.

Active ingredient: Metabolites of Trichoderma spp.

Treatment: Row Crops & Vegetables (Full Cover Foliar) Dosage: 0.5 ℓ per hectare Instructions: Add 0.5 ℓ MetaBOOST™ to 75 to 100 ℓ of unchlorinated water. Apply MetaBOOST™ as a foliar spray.

Treatment: Tree crops & Vines (Foliar) Dosage: 1 to 2 ℓ per hectare Instructions: Add 1 to 2 ℓ MetaBOOST™ to 100 ℓ of unchlorinated water. Apply MetaBOOST™ as a foliar spray.



39 JONAL DRIVE, CAVAN SOUTH AUSTRALIA 5094 PH: (08) 8359 6855

ADMINSA@WILCHEM.COM.AU WWW.WILCHEM.COM.AU

ABN 44 614 126 573

Manufactured by: Microbika (Pty) Ltd. Co. Reg No. 2015/194786/07 35 Eighth Avenu, Lichtenburg, 2740 South Africa



DIRECTIONS FOR USE: USE ONLY AS INDICATED

Compatibility:

Do NOT mix with any agrochemicals without consulting with your chemical distributor

Important notes:

Contents may settle. Shake well before use. Keep in a cool (4°C to 24°C) place. Do not allow to freeze. Store out of direct sunlight. Do not use product after expiry date.

Warnings and precautions:

Keep out of reach of children, uniformed persons and animals. May cause slight eye irritation. May cause allergic reactions in sensitive persons. Wash hands with soap and water before eating, drinking or smoking. Prevent contamination of food, feed, drinking water and eating utensils. Triple rinse and perforate empty container and do not use for any other purpose.

DV.	CE (NE	МА	NUFA	CTI	IDE:
UΗ	16 '	UT.	IVIAI	NUFA	UIL	JNE.

BATCH NO: