



SENTINEL RANGE

SENTINEL COPPER

EDTA Chelate

155 g/Kg Copper : 68 g/Kg Nitrogen
Wettable Micro Granule (WG)



Crop Type	Application	Rate Kg/Ha	Timing	Water (L/Ha)
Cereals Cotton, Maize, Summer Crops	Foliar	0.1 to 0.6	Apply from mid tillering to flag leaf emergence or when symptoms occur	75
	Soil	0.75 to 1		
Canola	Foliar	0.1 to 0.6	Apply at 4 to 8 leaf stage	75
Lucerne, Legumes, Pasture	Foliar	0.6 to 0.75	10 to 14 days prior to flowering	75
Vegetables	Foliar	0.1 to 0.2	Apply when sufficient foliage is present	500
	Fertigation	0.6 to 1.5		
Grapevines	Foliar	0.6 to 1	Apply when deficiency appears. * Fertigation rates are g/vine.	500
	Fertigation	2 to 6*		
Tree crops	Foliar	0.6 to 1	Apply early in the crop cycle as required. * Fertigation rates are g/tree.	500
	Fertigation	2 to 6**		

POISON - Keep out of reach of children.
FIRST AID - Avoid contact with eyes and skin. If poisoning occurs contact a doctor or poisons information centre on 131 120.

Any recommendations are a guide and must be adapted to suit local conditions. Adjust rates based on tissue tests, soil tests, local knowledge and the recommendation of your agronomist. Do not exceed a solution of 0.1% on fruit crops and 1% on broadacre crops.

Wilchem's Sentinel Range has been formulated to protect micronutrients in solution to allow for the correction of nutrient deficiencies.

Compatibility: It is compatible with a wide range of crop protection products. It is stable in alkaline soils and in the presence of soluble phosphates. A jar test should be performed and/or test application before preparing a full tank mix.

Directions for use: Do not apply during periods of crop stress, extreme weather, flowering or fruit colour change. Ensure all other products are mixed in the tank before adding. Do not apply as foliar to stone fruit.

Pack Size

20KG

Batch Number

SA Tel: (08) 8359 6855
SA Email: adminsa@wilchem.com.au

www.wilchem.com.au
ABN 44 614 126 573

WA Tel: (08) 9241 1188
WA Email: adminwa@wilchem.com.au