

Сгор Туре	Application	Rate Kg/Ha	Timing	Water (L/Ha)
Cereals Cotton, Maize, Summer Crops	Foliar	0.26 to 1	Apply from early tillering through to booting or when symptoms occur	60 - 80
	Soll	1 to 3		
Canola	Foliar	0.25 to 1	Apply at boiling	60 to 80
Lucerne and Legumes, Pasture	Foliar	0.25 to 1	10 to 14 days prior to flowering	60 to 80
Vegetables	Foliar	0.5 to 1.5	Apply as needed from 3 weeks after crop emergence	600
	Fertigation	3 to 7		
Grapevines	Foliar	0.6 to 1	Apply 2 to 3 time per season, flower bud visible and flower bud separated	260
	Fertigation	2 to 3		
Tree crops	Foliar	0.6 to 1	Apply as required. Othrus apply at bud burst. Apples apply at fruit set.	600
	Fertigation	2 to 3		

POISON - Keep out of reach of children.

FIRST AID - Avoid contact with eyes and skin. If poisoning occurs contact a doctor or poisons information centre on 131 126.

Any recommendations are a guide and must be adapted to sult local conditions. Adjust rates based on tissue tests, soil tests, local knowledge and the recommendation of your agronomist. Do not exceed a solution of 0.1% on fruit crops and 1% on broadacre crops.

Wilchem's Sentinel Range has been formulated to protect micronutrients in solution to allow for the correction of nutrient deficiencies.

Compatibility: It is compatible with a wide range of crop protection products. It is stable in alkaline soils and in the presence of soluble phosphates. A jar test should be performed and/or test application before preparing a full tank mix.

Directions for use: Do not apply during periods of crop stress, extreme weather, flowering or fruit colour change. Ensure all other products are mixed in the tank before adding. Do not apply as foliar to stone fruit.

Pack Size 20KG

Batch Number